ASPI | Afarand Scholarly Publishing Institute; Turkey
eISSN: 2645-7687. Journal of Clinical Care and Skills. 2025;6(4):219-223 10.58209/jccs.6.4.219

Red Blood Cell Alloimmunization in Thalassemia Patients in South-West Iran

ARTICLE INFO

Article Type
Descriptive Study

Authors
Rad F.*[1] *PhD*
Hadinia A.[1] *PhD*
Masoudifar S.[2] *MSc*

How to cite this article
Rad F, Hadinia A, Masoudifar S. Red Blood Cell Alloimmunization in Thalassemia Patients in South-West Iran. Journal of Clinical Care and Skills. 2025;6(4):219-223.

[1]Cellular and Molecular Research Center, Yasuj University of Medical Sciences, Yasuj, Iran
[2]Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tehran, Iran

Correspondence
Address: Cellular and Molecular Research Center, Yasuj University of Medical Sciences, Shahid Motahari Boulevard, Yasuj, Iran.
Postal Code: 7591994799
Phone: +98 (919) 6028093
fariba.rad89@gmail.com

Article History
Received: August 19, 2025
Accepted: September 29, 2025
ePublished: Octber 06, 2025

ABSTRACT

Aims Despite numerous studies on the prevalence of alloimmunization, its exact rate remains uncertain in many populations. In light of the Iranian Blood Transfusion Organization's recent national program to reduce alloantibody formation, this study investigated the prevalence of alloimmunization among thalassemia patients in Kohgiluyeh and Boyer-Ahmad Province, Iran, and explored associated factors, such as donor-recipient antigen homogeneity and early transfusion initiation.
Instrument & Methods This cross-sectional study was conducted on 134 beta-thalassemia major patients registered at Shahid Beheshti Hospital, Yasuj, Iran. Antibody screening was performed using both the standard tube method and the gel method in the immunohematology laboratory of Yasuj University of Medical Sciences.
Findings Out of 134 cases, 73 (54.48%) were female and 61 (45.52%) were male, with a mean age of 19.51±8.50 years. A high donor-recipient antigen homogeneity is suggested by the fact that 132 (98.5%) patients were native to Yasuj. All patients received ABO and Rh (D)-matched, leukodepleted blood transfusions at intervals ranging from 14 to 60 days (mean: 22.7±6.3 days); 65.7% were splenectomized. Alloantibodies were detected in two patients (1.49% prevalence; one with anti-K and one with anti-C), both of whom had undergone splenectomy. No autoantibodies or positive direct antiglobulin test results were observed.
Conclusion There is a low prevalence of RBC alloimmunization among BTM patients in Kohgiluyeh and Boyer-Ahmad Province.

Keywords Beta-Thalassemia; Blood Transfusion; Splenectomy

CITATION LINKS

[1] Anemia in children: With splenomegaly and peculiar changes in the bones ... [2] β-thalassemia distribution in the old world: An ancient disease seen from ... [3] Molecular basis of β thalassemia and potential ... [4] Epidemiology of thalassemia in Gulf Cooperation Council countries ... [5] Epidemiological investigation of a twenty-year major β-thalassemia surveillance ... [6] The frequency of beta-thalassemia mutations among carriers in Dezful ... [7] Thalassemia 2016: Modern medicine battles an ... [8] Global burden and unmet needs in the treatment of transfusion ... [9] Red blood cell transfusion risks in patients with end-stage ... [10] Red cell alloimmunization in repeatedly transfused sudanese patients with leukemia ... [11] Technical ... [12] Red blood cell alloimmunization in Iranian beta-thalassemia patients: A systematic review ... [13] Prevalence of alloimmunization against RBC antigens in thalassemia major patients of Lorestan ... [14] RBC alloimmunization in blood transfusion-dependent β-thalassemia patients ... [15] Prevalence of alloimmunisation in patients with beta ... [16] Determination of clinical prevalence and predominant pattern of RBCs alloimmunization among transfusion dependent thalassemic ... [17] Prevalence and specificities of red cell alloantibodies in transfusion-dependent beta thalassemia ... [18] Red cell alloimmunisation among sickle cell disease and thalassemia patients following rh-and k-matched red cell transfusion in Southwestern Saudi Arabia ... [19] Developing a databank for multiple transfusion patients: Rh antigen and phenotype distribution among 3000 regular blood ... [20] Prevalence of alloimmunization events in thalassemia patients with repeated transfusions in the rhesus blood group system: A systematic ... [21] From donor to recipient: Current questions relating to humoral ... [22] Splenectomy protects humans from red ... [23] Red blood cell alloimmunisation: Induction of immunity and potential ... [24] Human leukocyte antigen alloimmunization prevention mechanisms ... [25] An overview of red blood cell and platelet alloimmunisation ... [26] HLA polymorphisms and risk of red blood cell alloimmunisation in polytransfused ...

Copyright© 2025, the Authors | Publishing Rights, ASPI. This open-access article is published under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License which permits Share (copy and redistribute the material in any medium or format) and Adapt (remix, transform, and build upon the material) under the Attribution-NonCommercial terms.

Introduction

Thalassemia, first described by Thomas Cooley in 1925, is a genetic disorder caused by mutations in the globin genes, resulting in impaired hemoglobin synthesis that leads to defective erythropoiesis and severe anemia [1]. The term "thalassemia belt" refers to regions such as the Mediterranean, Southwestern Europe, Southeast Asia, and parts of Central Africa [2-4]. In Iran, particularly in areas surrounding the Caspian Sea and the Persian Gulf, thalassemia is hyperendemic [5, 6].

Regular blood transfusion remains the lifesaving treatment for thalassemia patients; however, it exposes them to adverse outcomes, including infections, hemosiderosis, and alloimmunization [7-9]. With each transfusion, a large number of donor-derived antigens are introduced into the recipient's circulation, potentially stimulating the immune system to produce alloantibodies against donor red blood cell (RBC) antigens that are absent on the recipient's RBCs—a phenomenon known as alloimmunization. Despite advances in pre-transfusion screening and extended phenotyping, alloimmunization continues to be a significant clinical challenge and may lead to delayed hemolytic reactions and difficulties in finding compatible blood units [10]. Few studies have explored alloimmunization specifically in Iran. The incidence of alloantibody formation in the thalassemia population ranges from 5.6% to 24.7% worldwide [11].

There are various studies on the rate of alloimmunization and the frequency of alloantibodies due to their importance in blood transfusion. According to a meta-analysis, the rate of alloimmunization is approximately 10% in Iran (95% CI: 7-13) [12]. One study on patients with thalassemia major in Lorestan Province, Iran, reports an alloimmunization rate of 1.53% [13]. In another study conducted in southern Iran, the prevalence of alloantibodies is 1.7% [14]. In some studies, the alloimmunization rate in beta-thalassemia major (BTM) patients is higher than that in our study. In the study by Kosaryan *et al.*, conducted in Sari, Iran, 40.4% of patients are identified as alloantibody positive using the tube method [15]. In another study by Hiradfar *et al.* in Ahwaz, Iran, 18.75% of patients are alloimmunized [16]. In similar studies conducted in different parts of Iran, the prevalence of alloimmunization has been reported to range from 2.87% (Northeast) to 40.4% (North) [17].

Despite these investigations, there are still regions in Iran with limited or no data on the prevalence of alloimmunization. Understanding these regional variations is essential for optimizing transfusion safety and guiding national blood policies. Therefore, the present study aimed to determine the prevalence of red cell alloimmunization among thalassemia patients in Kohgiluyeh and Boyer-Ahmad Province, Iran, and to evaluate potential influencing factors, such as age at first transfusion and splenectomy status.

Instrument and Methods

This cross-sectional study was conducted on 134 patients with BTM registered at Shahid Beheshti Hospital in Yasuj, Iran, between 2020 and 2021. Given the limited number of BTM patients in Yasuj (n=134) and considering that almost all of them regularly visit Cooley's Ward at Shahid Beheshti Hospital for scheduled blood transfusions every 15 to 45 days, the study population was defined as the entire reference population. Individuals with a confirmed diagnosis of BTM based on standard clinical and laboratory criteria and a regular transfusion history were included in the study. Patients were excluded if their medical records were incomplete or if they withdrew from participation during the study period.

After obtaining informed consent from each patient or from the patient's legal guardian (for minors), between 2 to 5mL of blood was collected from each patient in EDTA and clot tubes under standardized conditions. The EDTA samples were used for ABO and Rh (D) typing, as well as for the direct antiglobulin test (DAT), while the clot samples were used for antibody screening. Testing was conducted using Anti-A, Anti-B, and Anti-D reagents (Iranian Blood Research & Fractionation Company, Tehran, Iran) employing the standard tube method.

For the DAT, one drop of a 2-5% RBC suspension was added to four test tubes, followed by three saline washes. Polyspecific AHG, anti-IgG, and anti-C3 reagents (CE-Immun Diagnostika GmbH, Germany) were added simultaneously, with one tube serving as a negative control. After agitation and centrifugation, the tubes were examined for agglutination.

For the indirect antiglobulin test (IAT), two drops of patient serum were added to three separate tubes, followed by one drop each of screening cells I, II, and III (Biovesta, USA). After an immediate spin, albumin was added, and the samples were incubated at 37°C for 10 minutes. After washing, one drop of AHG was added to each tube; agglutination was then evaluated, and any negative reactions were rechecked using Coombs Control Cells (Biovesta, USA). Discrepant results between clinical findings and tube method results were confirmed with the gel method.

SPSS version 20 was used for data analysis utilizing the Chi-square test for the statistical analysis of qualitative data.

Findings

A total of 134 BTM patients were enrolled, of whom 73 (54.48%) were female and 61 (45.52%) were male. The mean age was 19.51±8.50 years (range: 1 to 37 years). Notably, 132 patients (98.5%) were

native to Yasuj, suggesting a high degree of donor-recipient antigen homogeneity.

The blood group distribution was as follows: O (48.5%), B (15.7%), A (30.6%), and AB (5.2%), with 97.8% positivity for the Rh (D) antigen. All patients received ABO and Rh (D)-matched, leukodepleted blood transfusions, which is a process where white blood cells (leukocytes) are removed from blood components. This is typically done using filters or centrifugation to reduce the number of leukocytes to a low level, with transfusions occurring at intervals of 14 to 60 days (mean interval: 22.7±6.3 days). Additionally, 88 patients (65.7%) underwent splenectomy. Splenectomy is a surgical procedure in which the spleen, an organ located in the upper left abdomen, is removed. It is often performed when the spleen is damaged, diseased, or enlarged, or when it is not functioning properly. The transfusion volume varied, with five patients receiving over 40 units annually and one patient receiving eight units within a year. The first transfusion was administered at a mean age of 15.42±19.40 months (range: 1 to 108 months).

Alloantibodies were detected in 2 patients (1.49% prevalence): one with anti-K and one with anti-C. Both of these patients had a history of splenectomy. No autoantibodies were observed using the standard tube method, and all DAT results were negative.

Discussion

This study aimed to determine the prevalence of red cell alloimmunization among BTM patients in Kohgiluyeh and Boyer-Ahmad Province, Iran. The prevalence of RBC alloimmunization among BTM patients in Kohgiluyeh and Boyer-Ahmad Province was found to be 1.49%, which is notably lower than reports from other regions in Iran, where the prevalence ranges from 0% to over 40% [12-17]. Understanding the factors contributing to this low alloimmunization rate is essential for developing preventive strategies and optimizing transfusion protocols. Several possible explanations for this low prevalence include high donor-recipient antigen homogeneity, early initiation of transfusions, leukodepleted blood transfusions, and splenectomy status [18, 19].

Studies conducted in different Iranian provinces have reported varying alloimmunization rates, influenced by genetic, geographical, and transfusion-related factors [20, 21]. As mentioned, a meta-analysis by Darvishi *et al.* estimates the overall alloimmunization rate in Iran to be approximately 10% (95% CI: 7-13) [12], which is considerably higher than the rate observed in our study. In Lorestan and southern Iran, alloimmunization rates are 1.53% and 1.7%, respectively [13, 14], aligning closely with our findings. However, studies in Sari and Ahvaz have reported significantly higher alloimmunization rates of 40.4% and 18.75%, respectively [15, 16]. The disparity may be attributed to differences in donor-recipient genetic compatibility, transfusion protocols, and variability in immune response.

Globally, alloimmunization rates among transfusion-dependent patients fluctuate due to regional diversity in blood donor genetics. In a large cohort from Saudi Arabia, the overall prevalence of RBC alloimmunization is 5.7%, with anti-K and anti-E being the most common alloantibodies [18]. In India, where thalassemia patients receive frequent transfusions, 15-20% develop alloantibodies, highlighting regional differences in transfusion practices and donor selection [19]. A study in the United States found that 30% of sickle cell disease (SCD) patients develop alloantibodies, largely due to racial differences between donors and recipients [20]. These findings suggest that ethnic background and antigen homogeneity between donors and recipients play a significant role in minimizing immune sensitization, particularly among BTM patients in homogeneous populations like Yasuj.

One of the strongest factors contributing to the low alloimmunization rate in our study is likely donor-recipient antigen homogeneity. Given that 98.5% of patients were native to Yasuj, their red cell antigen profiles likely closely resemble those of local blood donors, resulting in a lower likelihood of alloantibody formation. Antigen similarity between donors and recipients significantly reduces alloimmune responses [18, 19]. Studies have shown that alloimmunization rates are higher in regions with diverse ethnic backgrounds, where donor pools exhibit greater antigenic variation [20]. Thus, regional transfusion centers might benefit from HLA and extended RBC antigen matching programs to further minimize alloimmunization risks.

Another possible explanation for the low prevalence of alloimmunization is the early initiation of transfusions in BTM patients. Eighty-five percent of patients in this study had received their first transfusion before two years of age. Early transfusion is believed to induce immune tolerance, as the immune system is still developing and may not mount a strong alloimmune response [20, 21]. Studies have demonstrated that initiating transfusions early leads to lower alloimmunization rates compared to patients who begin transfusion therapy later in life. Optimizing transfusion protocols for neonates and infants may be an effective strategy for preventing alloimmunization.

The role of splenectomy in RBC alloimmunization is controversial, with conflicting results in different studies. Some studies suggest that splenectomy decreases immune sensitization, as the spleen plays a role in filtering foreign antigens, and splenectomy strongly reduces red blood cell alloimmunization [22, 23]. Others believe that splenectomized patients are more likely to develop alloantibodies due to prolonged exposure to transfused RBCs without efficient antigen clearance [21]. In our study, both

alloimmunized patients had undergone splenectomy, raising the possibility that splenectomy increases alloimmunization risk. Given the low number of alloimmunized cases, more multicenter studies are needed to evaluate the impact of splenectomy on alloantibody formation.

Leukodepletion is an important strategy for reducing alloimmunization, as leukocytes contribute to immune stimulation and cytokine production [24]. In our study, all patients received leukodepleted blood transfusions, which may explain the low alloimmunization rate. Previous studies have confirmed that the filtration of white blood cells reduces HLA antigen exposure, thereby lowering the risk of alloimmunization [25]. Standardizing universal leukodepleted transfusions for BTM patients could further reduce immune sensitization.

Despite its valuable findings, this study has several limitations. The small sample size (n=134) is indicative, but larger studies are needed to verify the findings. The low number of alloimmunized cases (n=2) limits statistical comparisons. Additionally, as a single-center study, a multicenter investigation could enhance generalizability. The lack of multivariate analysis means that future studies should employ logistic regression models to identify independent risk factors.

Further research should focus on expanding patient cohorts to enhance statistical validity, conducting genetic profiling of donors and recipients, and investigating HLA polymorphisms as potential determinants of alloimmunization risk [26].

The study revealed a low prevalence (1.49%) of RBC alloimmunization among BTM patients in Kohgiluyeh and Boyer-Ahmad Province, likely due to donor-recipient antigen homogeneity and early transfusion initiation. However, further large-scale studies with advanced statistical modeling are necessary to explore the impact of splenectomy, genetic diversity, and transfusion strategies on alloimmunization risk. Implementing standardized leukodepleted transfusion protocols and refining neonatal transfusion guidelines could significantly reduce the alloimmunization burden in transfusion-dependent patients.

Conclusion

There is a low prevalence of RBC alloimmunization among BTM patients in Kohgiluyeh and Boyer-Ahmad Province.

Acknowledgments: None declared by the authors.

Ethical Permissions: Written informed consent has been obtained from the patient(s) to publish this paper.

Conflicts of Interests: The authors declared no conflicts of interests.

Authors' Contribution: Rad F (First Author), Introduction Writer/Methodologist/Main Researcher/Discussion Writer/Statistical Analyst (70%); Hadinia A (Second Author), Introduction Writer/Assistant Researcher/Discussion Writer (20%); Masoudifar S (Third Author), Introduction Writer/Assistant Researcher/Discussion Writer (10%)

Funding/Support: This research received no external funding.

References

1- Cooley TB, Witwer ER, Lee P. Anemia in children: With splenomegaly and peculiar changes in the bones report of cases. Am J Dis Child. 1927;34(3):347-63.

2- De Sanctis V, Kattamis C, Canatan D, Soliman AT, Elsedfy H, Karimi M, et al. β-thalassemia distribution in the old world: An ancient disease seen from a historical standpoint. Mediterr J Hematol Infect Dis. 2017;9(1):e2017018.

3- Thein SL. Molecular basis of β thalassemia and potential therapeutic targets. Blood Cells Mol Dis. 2018;70:54-65.

4- Abu-Shaheen A, Heena H, Nofal A, Abdelmoety DA, Almatary A, Alsheef M, et al. Epidemiology of thalassemia in Gulf Cooperation Council countries: A systematic review. Biomed Res Int. 2020;2020(1):1509501.

5- Rezabeigi Davarani E, Mohseni Takaloo F, Vahidnia A, Daneshi S, Rezabeigi Davarani M, Khanjani N, et al. Epidemiological investigation of a twenty-year major β-thalassemia surveillance in Kerman, Iran. Arch Hyg Sci. 2020;9(4):265-74.

6- Joola P, Andashti B, Hosseini SA, Zadeh SMM, Bahrami N. The frequency of beta-thalassemia mutations among carriers in Dezful city, southwest Iran. Iran J Public Health. 2020;49(12):2438.

7- Rund D. Thalassemia 2016: Modern medicine battles an ancient disease. Am J Hematol. 2016;91(1):15-21.

8- Forni GL, Grazzini G, Boudreaux J, Agostini V, Omert L. Global burden and unmet needs in the treatment of transfusion-dependent β-thalassemia. Front Hematol. 2023;2:1187681.

9- Tanhehco YC, Berns JS. Red blood cell transfusion risks in patients with end-stage renal disease. Semin Dial. 2012;25(5):539-44.

10- Jawish TAA, Hessen HM, Alshafeea MA, Mohamedahmed KA, Ahmed EA, Modawe GA, et al. Red cell alloimmunization in repeatedly transfused sudanese patients with leukemia in Northern Sudan. Asian Pac J Cancer Prev. 2023;24(1):21-4.

11- Cohn CS, Delaney M, Johnson S, Katz LM, Schwartz J. Technical manual. Bethesda: Amer Assn of Blood Banks; 2023.

12- Darvishi P, Azami M, Sayehmiri K, Sayehmiri F, Goodarzi A, Azarkeivan A, et al. Red blood cell alloimmunization in Iranian beta-thalassemia patients: A systematic review and meta-analysis. ISBT Sci Ser. 2016;11(3):163-73.

13- Kiani A, Abdi J, Shirkhani Y, Kashi M. Prevalence of alloimmunization against RBC antigens in thalassemia major patients of Lorestan province in 1383. J Iran Blood Transfus. 2006;3(3):265-71. [Persian]

14- Karimi M, Nikrooz P, Kashef S, Jamalian N, Davatolhagh Z. RBC alloimmunization in blood transfusion-dependent β-thalassemia patients in southern Iran. Int J Lab Hematol. 2007;29(5):321-6.

15- Kosaryan M, Mahdavi MR, Roshan P, Hojjati MT. Prevalence of alloimmunisation in patients with beta thalassaemia major. Blood Transfus. 2012;10(3):396-7.

16- Hiradfar AA, Keikhai B, Pedram M. Determination of clinical prevalence and predominant pattern of RBCs alloimmunization among transfusion dependent

thalassemic patients in Ahvaz. Jundishapur Sci Med J. 2010;9(5):441. [Persian]
17- Vaziri M, JavadzadehShahshahani H, Moghaddam M, Taghvaee N. Prevalence and specificities of red cell alloantibodies in transfusion-dependent beta thalassemia patients in Yazd. Iran J Pediatr Hematol Oncol. 2015;5(2):93-9.
18- Meshi AA, Abu-Tawil H, Hamzi AA, Madkhali BA, Maghfori AB, Alnami II, et al. Red cell alloimmunisation among sickle cell disease and thalassemia patients following rh-and k-matched red cell transfusion in Southwestern Saudi Arabia: A multicenter study. Int J Gen Med. 2024;17:2855-64.
19- Bakhshandeh Z, Amirizadeh N, Maghsoodlu M, Oodi A, Naghi A, Khazaeli AA, et al. Developing a databank for multiple transfusion patients: Rh antigen and phenotype distribution among 3000 regular blood donors in Iran. Transfus Apher Sci. 2021;60(3):103124.
20- Indriani V, Mulyono B, Triyono T, Handayaningsih AE, Chandra LA. Prevalence of alloimmunization events in thalassemia patients with repeated transfusions in the rhesus blood group system: A systematic review and meta analysis. J Clin Med Res. 2025;17(2):106-18.
21- Prigent A, Maillard N, Absi L, Aloui C, Cognasse F, Laradi S, et al. From donor to recipient: Current questions relating to humoral alloimmunization. Antibodies. 2014;3(1):130-52.
22- Evers D, Van Der Bom JG, Tijmensen J, De Haas M, Middelburg RA, De Vooght K, et al. Splenectomy protects humans from red cell alloimmunization. Blood. 2016;128(22):24.
23- Hendrickson JE. Red blood cell alloimmunisation: Induction of immunity and potential mitigation strategies. ISBT Sci Ser. 2018;13(1):105-11.
24- Adane T, Enawgaw B. Human leukocyte antigen alloimmunization prevention mechanisms in blood transfusion. Asian J Transfus Sci. 2023;17(2):264-72.
25- Garraud O, Chiaroni J. An overview of red blood cell and platelet alloimmunisation in transfusion. Transfus Clin Biol. 2022;29(4):297-306.
26- Rodrigues C, Sell AM, Guelsin GAS, Higa TT, Pagliarini E Silva S, Macedo LC, et al. HLA polymorphisms and risk of red blood cell alloimmunisation in polytransfused patients with sickle cell anaemia. Transfus Med. 2017;27(6):437-43.